



# Allergologia et immunopathologia

Sociedad Española de Inmunología Clínica,  
Alergología y Asma Pediátrica

[www.elsevier.es/ai](http://www.elsevier.es/ai)



## ORIGINAL ARTICLE

# Cross-over clinical trial for evaluating the safety of camel's milk intake in patients who are allergic to cow's milk protein



E.M. Navarrete-Rodríguez<sup>a</sup>, L.A. Ríos-Villalobos<sup>a</sup>, C.R. Alcocer-Arreguín<sup>a</sup>,  
B.E. Del-Rio-Navarro<sup>a,\*</sup>, J.M. Del Rio-Chivardi<sup>a</sup>, O.J. Saucedo-Ramírez<sup>a</sup>,  
J.J.L. Sienna-Monge<sup>b</sup>, R.V. Frias<sup>c</sup>

<sup>a</sup> Department of Allergy and Clinical Immunology, Hospital Infantil de México Federico Gómez, Ciudad de México, Mexico

<sup>b</sup> Department of Ambulatory Pediatrics, Hospital Infantil de México Federico Gómez, Ciudad de México, Mexico

<sup>c</sup> Department of Gastroenterology and Nutrition, Hospital Infantil de México Federico Gómez, Ciudad de México, Mexico

Received 24 March 2017; accepted 21 June 2017

Available online 6 December 2017

### KEYWORDS

Cow's milk protein allergy (CMPA);  
Camel milk

### Abstract

**Background:** Cow's milk protein allergy (CMPA) affects between 0.6 and 0.9% of the general population, and its treatment implies the total elimination of the intake of this protein. Camel's milk has been suggested as an alternative for patients over one year of age who suffer from CMPA due to the difference in the amino acid sequence from that of cow's milk. The objective of this study was to evaluate the safety and tolerability of camel's milk in children with CMPA.

**Methods:** Crossed clinical trial for the use of camel's milk vs. amino acid formula, carried out at the Dr. Federico Gómez Children's Hospital of Mexico (HIMFG) on patients between one and 18 years of age with diagnosed CMPA confirmed through double-blind, placebo-controlled food challenges (DBPCFCs). Only those whose allergies were confirmed were randomly placed into two groups: those to be administered camel's milk and those to be administered the amino-acid formula for two weeks, followed by a six-week wash-out period, and then a group crossing for a further two weeks.

**Results:** 49 patients with suspected CMPA were included in the study; the diagnosis was confirmed through DBPCFCs in 15 patients, who were those who participated in the study. After having been administered camel's milk, none of the patients presented adverse effects.

**Conclusions and clinical relevance:** Camel's milk is safe and tolerable in patients above one year of age with CMPA and can be considered as a good alternative given the benefit of its taste compared to other formulas.

© 2017 SEICAP. Published by Elsevier España, S.L.U. All rights reserved.

**Abbreviations:** CMPA, cow's milk protein allergy; HIMFG, Hospital Infantil de México Federico Gómez; DBPCFC, double-blind, placebo-controlled food challenge.

\* Corresponding author.

E-mail address: [blancadelrionavarro@gmail.com](mailto:blancadelrionavarro@gmail.com) (B.E. Del-Rio-Navarro).

<https://doi.org/10.1016/j.aller.2017.06.005>

0301-0546/© 2017 SEICAP. Published by Elsevier España, S.L.U. All rights reserved.

## Introduction

Cow's milk protein allergy (CMPA) affects between 0.6 and 0.9% of the general population, and is a public health issue that mainly affects the paediatric population.<sup>1,2</sup> Although the diagnosis of CMPA is not simple, we know that double-blind, placebo-controlled food challenges (DBPCFCs) continue to be the gold standard for its diagnosis, especially in research studies.<sup>2-4</sup> Current treatment for CMPA today implies totally eliminating the intake of the protein, which in most cases represents a huge challenge for the family and attending physicians.<sup>5-7</sup>

Whenever prescribing a diet free from cow's milk protein, it must be nutritionally adequate, which is why the objective is to reach a protein-calorie balance with an adequate amino-acid composition, as well as a proper source of calcium. It is necessary to complement the diet of children under the age of one with an extensively hydrolysed or amino-acid formula,<sup>8</sup> however, low organoleptic acceptance, understood as the sensory perception of the physical qualities of matter, greatly limits the use of these formulas, as does adherence; furthermore, it has been noted that the smaller the amount of administered protein, the worse the flavour.<sup>9</sup>

After the first year of age, introducing solid foods into the baby's diet avoids the need for cow's milk protein.<sup>8</sup> Formulas that are extensively hydrolysed with casein and whey, as well as amino acids, are designed for children under the age of one, acting as a substitute for cow's milk protein; after this first year of life, this protein is no longer necessary for their diet.<sup>8</sup> However, it has been proven that children over the age of one who have restrictions regarding this product and its derivatives, absorb less protein and calories over all, even when aided by nutritionists.<sup>10</sup> Cow's milk is highly consumed not only in Mexico, but around the world. According to the Mexican Secretariat of Economy and the General Directorate for Basic Industries, in 2010, 97 litres of cow's milk were consumed per inhabitant per year in Mexico,<sup>11</sup> without taking into account that a significant amount of processed food contains cow's milk protein as a base ingredient.<sup>12,13</sup>

Camel's milk has been proposed as an alternative for patients who suffer from CMPA because of its different amino-acid sequence; furthermore, it presents low quantities of  $\beta$  casein (Bos d8) and no  $\beta$  lactoglobulins (Bos d5),<sup>14</sup> and is adequately nutritionally balanced with essential components such as lactoferrin, immunoglobulins, lysozyme, and vitamin C<sup>15</sup> (Table 1).

Various authors have tried to test the tolerability of camel's milk on patients who are allergic to cow's milk protein, and have obtained favourable results; however, it is worth considering that most of these studies evaluate the tolerability of camel's milk in open trials, which could skew the results.

The objective of our study was to evaluate the safety of administering camel's milk to paediatric patients over the age of one with CMPA.

## Materials and methods

A cross-over clinical trial was carried out at the Dr. Federico Gómez Children's Hospital of Mexico (HIMFG), from

January to December 2016, to compare the safety between administering camel's milk vs. amino-acid formula on children diagnosed with CMPA. Patients from 1 to 18 years of age with suspected CMPA – understood as the presence of respiratory, cutaneous, gastrointestinal or nasal symptoms associated with the intake of cow's milk protein – were included in the initial study. Patients with allergies from other food sources were excluded from the study, as well as those under the treatment of immunosuppressive drugs, systemic steroids, antihistamines and anti-leukotrienes. None of the patients suffered from any concomitant chronic illnesses, except from rhinitis or asthma; notwithstanding, adequate symptom control was carried out when participating in the trial.

This trial was approved by the Ethics and Biosafety investigation committees at HIMFG (protocol number HIM/2016/006).

All patients, with prior informed consent, underwent a full medical evaluation by a certified paediatrician, including medical records and a physical examination. The first step was to confirm the diagnosis of CMPA by carrying out a DBPCFC with the cow's milk protein; in accordance with international guidelines on the matter,<sup>23</sup> patients underwent a strict cow's milk protein elimination diet over six weeks with previous supervision and education regarding food that could contain the ingredient and label reading. At the end of these six weeks, a placebo or lactose-free milk was administered on two separate days, with dosage levels progressing every 30 min, until reaching a volume of 240 ml. The presence of abdominal, cutaneous, respiratory and nasal symptoms was monitored, and results were considered positive when objective symptoms were present, or subjective symptoms in three consecutive doses, and worsened or persisted after 40 min.<sup>24</sup>

Having confirmed the diagnosis of CMPA, medical advice was newly given concerning the elimination diet and patients were summoned six weeks later. During this six-week period, patients were randomised by a random numbers generator which balanced out segments of five patients in one of two groups: one to be administered camel's milk and the other an amino-acid formula in accordance with Table 2. Neither the doctors in charge of administering and monitoring, nor the patients nor family members knew what the trial was about. Patient monitoring was carried out according to the times specified in Table 2 at the hospital, in case of the presence of abdominal, cutaneous, respiratory and nasal symptoms. Results were deemed positive when objective symptoms were present, or subjective symptoms present in three consecutive doses, and worsened or persisted after 40 min.

In case of negative results throughout this period of hospital monitoring, patients would be released and given enough formula/milk so as to ingest 200 ml per day over seven days. At the end of the intervention, an intermediate six-week cleansing period was given, in which neither intervention (formula or milk) was administered. The groups were then switched, so as to carry out the same procedure with the other intervention. Daily follow-up was given over the phone during these two weeks, and at the end of the whole process a physical examination was carried out by a paediatric allergist.

**Table 1** Nutritional composition of 100 ml of camel milk and other milk formulas.<sup>25–27</sup>

Components	Camel milk	Cow milk	Human milk	Extensively hydrolysed formula	Amino-acid formula
Energy (kcal)	45	62.7	62	75	75
Proteins (g)	2.2	3.2	0.9	2.0	2
Fat (g)	2	3.5	3.2–4.5	3.7	3.7
Whey/casein Relation	30:70	20:80	80:20	100:0 <sup>a</sup>	N/A
Carbohydrates (g)	4.6	4.6	6.8–7.4	8	8.5
Phosphorus (mg)	51.5	91	14	34.4	40
Magnesium (mg)	14.2	9.6	2.8	7.9	7
Calcium (mg)	117	115	28	51.3	60
Potassium (mg)	60	140	45	83.4	86.6
Chlorine (mg)	163	97	39	59.6	63
Sodium (mg)	69	51.5	14	33	26.3
Copper (mcg)	1.4	3	35	54	61.3
Iron (mcg)	39	0.05	76	0.6	0.76
Zinc (mcg)	121	0.4	240	0.6	0.76
Vitamin C (mg)	4.6	3	3.5	8.7	10
Thiamine (B <sub>1</sub> ) (mg)	0.05	0.03	0.01	0.47	0.07
Riboflavin (B <sub>2</sub> ) (mg)	0.07	0.09	0.04	0.09	0.16
Niacin (mg)	0.52	0.16	0.21	0.66	0.62

<sup>a</sup> The composition of milk formulas may vary depending on the manufacturer, for the determination of breast milk the mature stage was considered.

**Table 2** Steps for administration of double-blind placebo-controlled oral challenge.

Step	Time	Amount	Tolerates
1	00:00	0.1 ml	Yes <input type="checkbox"/> No <input type="checkbox"/>
2	00:30 min	0.2 ml	Yes <input type="checkbox"/> No <input type="checkbox"/>
3	01:00 h	0.5 ml	Yes <input type="checkbox"/> No <input type="checkbox"/>
4	01:30 h	1.5 ml	Yes <input type="checkbox"/> No <input type="checkbox"/>
5	02:00 h	4.5 ml	Yes <input type="checkbox"/> No <input type="checkbox"/>
6	02:30 h	15 ml	Yes <input type="checkbox"/> No <input type="checkbox"/>
7	03:00 h	40 ml	Yes <input type="checkbox"/> No <input type="checkbox"/>
8	03:30 h	60 ml	Yes <input type="checkbox"/> No <input type="checkbox"/>
9	04:40 h	97.2 ml	Yes <input type="checkbox"/> No <input type="checkbox"/>
Total	06:00 h	219 ml	

Verum preparation (camel milk): amino-acid formula, camel milk (2.62 g protein), sugar + vanilla flavouring.  
Placebo preparation (amino-acid formula): amino-acid formula, sugar, vanilla flavouring.

## Results

A total of 49 patients with suspected CMPA were evaluated, 49.0% male (95%CI 35.5–62.5) and 51.0% female (95%CI 37.4–64.4); at the time of trial inclusion, the mean age was 4.3 (1–13 years old). The most frequently mentioned

symptoms related to cow's milk protein consumption were abdominal and mainly abdominal pain (Table 3).

A DBPCFC for cow's milk protein was carried out on the 49 trial patients, confirming CMPA diagnosis in one third of the total (15 patients), nine males and six females with a mean age of four (1–9 years old) (Table 3).

The most frequent symptoms in patients with confirmed CMPA were cutaneous, then abdominal (Table 4).

The 15 patients with diagnosed CMPA were randomised to receive one of the two interventions (Fig. 1). All of the patients concluded the protocol. None of the 15 patients presented any symptoms from the amino-acid formula or from the camel's milk during the days the trial was carried out, hence they followed the free daily intake of 200 ml of camel's milk over the following seven days.

None of the patients presented late adverse effects from ingesting camel's milk.

## Discussion

The prevalence of CMPA is a significant public health issue that has increased over the last decades,<sup>2</sup> and whose treatment implies the total suspension of the intake of this protein. The clinical use of milk originating from other mammals has been proposed, including that of sheep, goats, donkeys and buffalo, however, their protein sequences have

**Table 3** Description of the population included in the study in groups: camel milk and amino-acid formula.

	Camel milk <i>n</i> = 8	Amino-acid formula <i>n</i> = 7	<i>p</i>
Sex	Male = 62.5% Female = 37.5%	Male = 57.1% Female = 42.8%	<i>p</i> = 0.62
Age	Median = 4.4 years old (1.5–7 years old)	Median = 2.9 years old (1.8–9.5 years old)	<i>p</i> = 1.00
Age at diagnosis	Median = 2.5 years old (1–7 years old)	Median = 1.8 years old (1–9.5 years old)	<i>p</i> = 1.00

**Table 4** Symptoms presented in patients with DBPCFC positive to cow's milk.


Present symptoms during DBPCFC to cow's milk	Percentage n = 15	95%CI
Abdominal	73.3%	48.0–89.1%
Skin	80.0%	54.8–92.9%
Respiratory	26.6%	10.9–51.9%
Cardiovascular	0%	NV
Anaphylaxis	0%	NV

DBPCFC were performed to 49 patients with suspected CMPA, the results presented here are those obtained in patients in whom the diagnosis was confirmed, for the remaining patients CMPA was discarded and they currently consume cow's milk protein without presenting any symptoms. NV: not valid.

been found to be homologous to that of cow's milk, which is why their use is not recommended in CMPA patients.<sup>2</sup> Camel's milk has been suggested as an alternative for this

population due to its differences from cow's milk protein<sup>19</sup> as well as its high nutritional content; camel's milk has been widely distributed and accepted in countries in East Africa for decades, and with the growing levels of urbanisation, consumption and commercialisation have increased in urban areas, improving quality standards, and managing sales in countries far from the Middle East.<sup>20–22</sup>

We did not find mediated or non-mediated IgE symptoms regarding the intake of camel's milk, proving a similar tolerability level between the amino-acid formula and camel's milk in CMPA patients, as found by El-Agamy et al.,<sup>16</sup> who administered camel's milk to a group of 28 children with confirmed CMPA without any secondary clinical manifestations in its use; also Shabo et al.,<sup>17</sup> who through a simple trial administered camel milk to a group of eight children with CMPA, proving adequate tolerance. Ehlal et al. studied a group of 35 patients with symptoms related to CMPA with specific IgE and positive skin tests for cow's milk; all patients underwent skin tests for camel's milk and those with



**Figure 1** Patient enrollment, randomization and follow-up.

negative results ( $n=28$ ) were administered camel's milk. None of those 28 patients presented any symptoms while undergoing the open trial.<sup>18</sup>

Throughout the ample literature review we carried out, we found only one reported case where a patient with CMPA suffered from anaphylaxis with camel's milk, published by Al-Hammadi<sup>23</sup> in 2010, concerning an atopic six-year-old child with uncontrolled asthma, multiple food allergies and atopic dermatitis; we considered it unwise to carry out DBPCFCs with camel's milk on patients with uncontrolled allergic pathologies, especially with asthma, due to the high risk of severe side effects.

It is worth noting that as suggested in worldwide literature, we found a significant overdiagnosis of CMPA, which was only confirmed in 30.6% of patients who suspected suffering from CMPA.

Having an alternative to cow's milk for children with CMPA, which has an adequate taste and is safe to consume, is extremely important, as it could have a positive impact on the quality of life of patients and their families. Up until now, our study is the only one to have been carried out with DBPCFCs to value the tolerability of camel's milk in patients with CMPA, significantly reducing methodological bias and adding value to the presented results.

Although other authors have performed skin tests or measured camel's milk's specific IgE, we did not carry out those tests because we used DBPCFC, which is the gold standard of diagnosis and capable of diagnosing both IgE and non-IgE CMPA.

## Conclusion

Our study proves that camel's milk is safe and adequately tolerable for patients over the age of one with CMPA and could be considered as a good alternative due to its better flavour and lower cost than other formulas.

## Funding

Camel's milk was obtained as a donation from the Emirate company "Camelicious" Emirates Industry for Camel Milk & Products.

## Conflict of interest

The authors declare no conflict of interest.

## Acknowledgments

The authors thank "Camelicious" Emirates Industry for Camel Milk & Products for supplying camel milk and to Muhammad Ashraf, Sales Manager, who was personally in charge of the process.

## References

- Chafen JJ, Newberry SJ, Riedl MA, Bravata DM, Maglione M, Suttortp MJ, et al. Diagnosing and managing common food allergies: a systematic review. *JAMA*. 2010;303:1848–56.
- Fiocchi A, Schünemann HJ, Brozek J, Restani P, Beyer K, Troncone R, et al. Diagnosis and rationale for action against cow's milk allergy (DRACMA). *J Allergy Clin Immunol*. 2010;126:1119–28.
- De Greef E, Hauser B, Devreker T, Veereman-Wauters G, Vandenas Y. Diagnosis and management of cows milk protein allergy in infants. *World J Pediatr*. 2012;8:19–24.
- Huang F, Kim JS. IgE-mediated cows milk allergy in children. *Curr Allergy Asthma Rep*. 2012;12:630–40.
- Dambacher WM, de Kort EH, Blom WM, Houben GF, de Vries E. Double-blind placebo-controlled food challenges in children with alleged cow's milk allergy: prevention of unnecessary elimination diets and determination of eliciting doses. *Nutr J*. 2013;12:22.
- Kattan J, Cocco R, Jarvinen K. Milk and Soy Allergy. *Pediatr Clin N Am*. 2011;58:407–26.
- Vandenas Y, Brueton M, Dupont C, David Hill, Erika Isolauri, Sibylle Koletzko, et al. Guidelines for the diagnosis and management of cows milk protein allergy in infants. *Arch Dis Child*. 2007;92:902–8.
- Koletzko S, Niggemann B, Arato A, Dias JA, Heuschkel R, Husby S, et al. Diagnostic approach and management of cow's-milk protein allergy in infants and children: ESPGHAN GI Committee practical guidelines. *J Pediatr Gastroenterol Nutr*. 2012;55:221–9.
- Pedrosa M, Pascual CY, Larco JI, Esteban MM. A comparative study of taste, smell, and texture evaluated by healthy volunteers. *J Invest Allergol Clin Immunol*. 2006;16:351–6.
- Mastin K, Oliver EM, Scally KS, Atkinson J, Foote K, Venter C, et al. Nutritional adequacy of a cows' milk exclusion diet in infancy. *Clin Transl Allergy*. 2016;6.
- [http://www.economia.gob.mx/files/comunidad\\_negocios/industria\\_comercio/informacionSectorial/analisis\\_sector\\_lacteo.pdf](http://www.economia.gob.mx/files/comunidad_negocios/industria_comercio/informacionSectorial/analisis_sector_lacteo.pdf).
- Robson SM, Khoury JC, Kalkwarf HJ, Copeland K. Dietary intake of children attending full-time child care: what are they eating away from the child-care center? *J Acad Nutr Diet*. 2015;115:1472–8.
- Meyer R, De Koker C, Dziubak R, Godwin H, Dominguez-Ortega G, Shah N. Dietary elimination of children with food protein induced gastrointestinal allergy – micronutrient adequacy with and without a hypoallergenic formula? *Clin Transl Allergy*. 2014;4:31.
- Ehlayel M, Bener A, Abu Hazeima K, Al-Mesaifri F. Camel milk is a safer choice than goat milk for feeding children with cow milk allergy. *ISRN Allergy*. 2011;29:391641.
- De Almeida Roberto Ronald. Leche de camello: características y perspectivas para uso en clínica. *Rev Chil Nutr*. 2011;38:211–8.
- El-Agamy El, Nawar M, Shamsia SM, Awad S, Haenlein GFW. Are camel milk proteins convenient to the nutrition of cow milk allergic children? *Small Rumin Res*. 2009;82:1–6.
- Shabo Y, Barzel R, Margoulis M, Yagil R. Camel milk for food allergies in children. *Isr Med Assoc J*. 2005;7:796–8.
- Ehlayel MS, Hazeima KA, Al-Mesaifri F, Bener A. Camel milk: an alternative for cow's milk allergy in children. *Allergy Asthma Proc*. 2011;32:255–8.
- Restani P, Beretta B, Fiocchi A, Ballabio C, Galli CL. Cross-reactivity between mammalian proteins. *Ann Allergy Asthma Immunol*. 2002;89:11–5.
- El-Agamy El. Are camel milk proteins convenient to the nutrition of cow milk allergic children? *Small Rumin Res*. 2009;82:1–6.
- Farah Z. Camel milk. *Swiss Federal of Technology*. Elsevier; 2011.
- Mohammad S, Ehlayel MD, Khalid Abu Hazeima MD. Camel milk: an alternative for cow's milk allergy in children. *Allergy Asthma Proc*. 2011;32:255–8.
- Al-Hammadi S, El Hassan T, Al-Reyami L. Anaphylaxis to camel milk in an atopic child. *Allergy*. 2010;65:1622–9.
- Sampson HA, van Wijk RG, Bindslev-Jensen C, Sicherer S, Teuber SS, Burks AW, et al. Standardizing double blind, placebo-

- controlled oral food challenges: American Academy of Allergy and Clinical Immunology PRACTALL consensus report. *J Allergy Clin Immunol.* 2012;130:1260–74.
25. Product Nutrition Labeling. <http://www.fao.org/docrep/003/X6528E/X6528E03.htm#chIII>. <https://camelicious.in/shop/camel-milk-powder/buy-camel-milk-india-1kg/>.
26. Fomon Samuel J. *Nutrition of normal infants*, vol. 1, 1st ed. Mosby; 1993 [chapters 9 and 10].
27. Shamsia SM. Nutritional and therapeutic properties of camel and human milks. *Int J Genet Mol Biol.* 2009;1:52–8.